




SCW Confidential Performance Report for Hospital X
Quality Initiative: Reducing Postoperative Opioid Prescribing
Reporting Period: January-December 2017

Figure 1. Unadjusted mean morphine equivalent dose provided to patients following laparoscopic cholecystectomy



Black line: Statewide mean morphine equivalent dose. Black dashed line: Target dose. Each bar represents a hospital in Wisconsin. Error bars represent confidence intervals around each hospital estimate. Your hospital’s performance is represented by the solid blue bar.

Figure 2. Case volume and opioid initial fill preceding or following laparoscopic cholecystectomy

	Opioid Prescribing Recommendation*	Hospital X	Participating Hospitals (n=)	All WI Hospitals (n=)
Number of Cases	n/a			
Hydrocodone (Norco) 5 mg Tablets (Mean, SD)	15			
Codeine (Tylenol #3) 30 mg Tablets (Mean, SD)	15			
Tramadol 50 mg Tablets (Mean, SD)	15			
Oxycodone 5 mg Tablets (Mean, SD)	10			
Hydromorphone (Dilaudid) 2 mg Tablets (Mean, SD)	10			
Payer	n/a			
Private insurance				
Medicare				
Medical assistance/Badgercare				
Other				

* Prescribing recommendation taken from the Michigan Opioid Prescribing Engagement Network (OPEN)

